



TC1 Test 01/01/1952

CLENPIQ Constipation AM - Colonoscopy Preparation Instructions

Appointment Date: Check-in Time: Procedure Performed by:

You will need:

- Your Clenpiq bowel prep. Please pick up ahead of time. This prescription was sent to your pharmacy when your procedure was scheduled.
- **Magnesium Citrate** - One 10oz. bottle (No Red). It is available in the laxative section of your pharmacy.

Special Medication and Diabetic Directions: See separate instructions.

One week before

Stop taking iron pills, vitamins or herbal supplements in the pill form. Take your usual prescription medication during prep unless told not to. Aspirin, Ibuprofen and Tylenol are OK if needed.

Avoid foods with seeds (flax seeds, popcorn, nuts and berries). They clog the colonoscope.

Two days before

Stop taking fiber supplements: Metamucil, Citrucel, etc.

Begin clear liquid diet. You may have: water, coffee, tea, (No milk or creamers) clear fruit juices without pulp or cloudiness, soda pop, Gatorade, chicken bouillon, popsicles, and JELL-O.

Avoid the color **red** in all liquids. Avoid alcohol.

5:00 pm: Drink the 10 oz. bottle of Magnesium Citrate (No Red) followed by a full glass of water.

1 Day before your colonoscopy

Morning: No Solid Food. Continue clear liquid diet.

NO SOLID FOOD. You may have: water, coffee, tea, (No milk or creamers) clear fruit juices without pulp or cloudiness, soda pop, Gatorade, chicken bouillon, popsicles, and JELL-O. Avoid the color red in all liquids. Avoid alcohol.

4 PM: Drink one bottle of Clenpiq. Drink Clenpiq straight from the bottle. *Do not refrigerate or freeze Clenpiq.* Stay extremely close to your restroom.

Over the next 5 hours drink 5 cups (8 oz each) of clear liquids.

11 PM: Drink one bottle of Clenpiq. Drink Clenpiq straight from the bottle. *Do not refrigerate or freeze Clenpiq.* Stay extremely close to your restroom.

Over the next 5 hours drink 3 cups (8 oz each) of clear liquids.

Day of your colonoscopy *DO NOT EAT OR DRINK WITHIN 2 HOURS OF YOUR PROCEDURE!*

Take your regular morning medications at least 4 hours before your appointment

DO NOT put ANYTHING in your mouth the 2 hours before your procedure.

This includes but is not limited to: bowel prep, medication, hard candy, chewing gum and chewing tobacco.

No Food. No Drink. 2 hours.

If these instructions have not been followed your appointment may be delayed or cancelled.

If you received these instructions by email, you will also receive an attachment with additional. important information.