

CLENPIQ Constipation PM - Colonoscopy Preparation Instructions

Appointment Date:

Check-In Time:

Procedure Performed by:

You will need:

- Your Clenpiq bowel prep. Please pick up ahead of time. This prescription was sent to your pharmacy when your procedure was scheduled. *Please contact our office if the prescription is not covered or has a high cost*
- **Miralax (polyethylene glycol)** - 34 grams - It is available over the counter in the laxative section.
- GasX or Simethicone gas relief capsules. Purchase over-the-counter. You will need 4 of these.

Special Medication and Diabetic Directions: See separate instructions.

One week before

Stop taking iron pills, vitamins or herbal supplements in the pill form.

Take your usual prescription medication during prep unless told not to. Aspirin, Ibuprofen and Tylenol are OK if needed.

Avoid foods with seeds (flax seeds, popcorn, nuts and berries). They clog the colonoscope.

Two days before

Stop taking fiber supplements: Metamucil, Citrucel, etc.

When you awake, begin Clear Liquid Diet. No Solid Food.: You may have: water, coffee, tea, (No milk or creamers) clear fruit juices without pulp or cloudiness, soda pop, Gatorade, chicken bouillon or clear broth, popsicles, and JELL-O.

Avoid the color **red** in all liquids. Avoid alcohol.

5:00 pm: Mix 34 grams of Miralax (polyethylene glycol) in 16 ounces of water and drink.

1 Day before your colonoscopy

Continue Clear Liquid Diet. No Solid Food.

6 PM - Drink one bottle of Clenpiq. Drink Clenpiq straight from the bottle. Do not refrigerate or freeze Clenpiq. Stay extremely close to your restroom.

Take 2 GasX pills.

Over the next 5 hours drink 5 cups (8 oz each) of clear liquids.

DO NOT EAT OR DRINK WITHIN 2 HOURS OF YOUR PROCEDURE!

Day of your colonoscopy

Continue Clear Liquid Diet. No Solid Food.

5-6 hours prior to your check in time: Drink one bottle of Clenpiq. Drink Clenpiq straight from the bottle. Do not refrigerate or freeze Clenpiq. Stay extremely close to your restroom.

Take 2 GasX pills.

Drink at least 3 cups (8 oz each) clear liquids. Finish clear liquids at least 2 hours before your procedure.

Take your regular morning medications at least 2 hours before your appointment. Please do not take any narcotics or controlled substances, which includes some anxiety medications until after your procedure.

DO NOT put ANYTHING in your mouth the 2 hours before your procedure.

This includes but is not limited to: bowel prep, medication, hard candy, chewing gum and chewing tobacco.

No Food. No Drink. 2 hours.

You also need to arrange for an **adult driver** that is able to walk on their own, help you in and out of a car, drive you home and remain with you for 6 hours.

If these instructions have not been followed your appointment may be delayed or cancelled.

If you received these instructions by email, you will also receive an attachment with additional, important information.